

COARSE-GRAINING AND CRITICAL SPIN CORRELATIONS IN PYROCHLORE ANTIFERROMAGNETS

C. L. Henley

Dept. of Physics, Clark Hall, Cornell Univ., Ithaca NY 14853-2501

Abstract

Consider the Ising spin antiferromagnet on the pyrochlore lattice consisting of corner-sharing tetrahedra. (Isotropic 3-component spins are a minor extension.) The ground states are highly degenerate and map to configurations of arrows on the edges of a diamond lattice; the ground-state constraint that every tetrahedron has zero net spin translates to the “ice rule” (two arrows in and two out of every diamond-lattice vertex), as is well known. Consequently, the vector field of coarse-grained arrows $\mathbf{E}(\mathbf{r})$ has zero divergence; furthermore, the free energy (from entropy) has the form $\int d^3\mathbf{r} |\mathbf{E}(\mathbf{r})|^2$. Thus the arrow field behaves as a pseudo “electric field” without sources. This has the following consequences:

(a) The spin correlations have the form of a dipole-dipole interaction, thus they are *critical* (decay with distance as a power-law). This is already known in the ice context [1, 2].

(b) The diffuse magnetic scattering has a very low intensity over the entire first Brillouin zone; it is virtually zero along the (100) axes, and exhibits pinch-point (pseudo-dipolar) singularities at (200) and (111) points. All this is observed in pyrochlore systems experimentally[3] and in simulations[4]. An extension of this approach produces a phenomenological fitting formula for diffuse intensity throughout the Brillouin zone, similar in form to Ref.[5].

(c) The AC susceptibility behavior[6] should be cast in terms of the diffusion and recombination of pseudo “electric charges”.

References

- [1] F. H. Stillinger and M. A. Cotter, J. Chem. Phys. 58, 2532 (1973); J. Villain and J. Schneider, in *Physics and Chemistry of Ice*, ed. E. Whalley *et al* (1973).
- [2] R. W. Youngblood and J. D. Axe, Phys. Rev. B 23, 232 (1981).
- [3] R. Ballou, E. Lelièvre-Berna, and B. Fåk, Phys. Rev. Lett. 76, 2125-8 (1996); S-H Lee, C. Broholm, W. Ratcliff, G. Gasparović, Q. Huang, T. H. Kim, and S.-W. Cheong, Nature 418, 856 (2002).
- [4] M. J. Harris, M. P. Zinkin, and T. Zeiske, Phys. Rev. B 56, 11786-90 (1997); B. Canals and C. Lacroix, Phys. Rev. Lett. 80, 2933-6 (1998).
- [5] J.N.Reimers, Phys.Rev.B 46, 193 - 202 (1992).
- [6] J. Snyder, J.S. Slusky, R.J. Cava, and P. Schiffer, Phys. Rev. B 66, 064432 (2002).